

# Compact Sets

MATH 409 HNR Analysis on the Real Line, Texas A&M University

Joshua Im

April 29, 2025

Recall the definition of compact sets discussed:

## Definition 0.1: Compact Sets

A set  $S \subset \mathbb{R}$  is **compact** if every sequence in  $S$  has a convergent subsequence with limit in  $S$ .

This definition is enough for  $\mathbb{R}$ , but there is a generalized definition.

1

## Metric Spaces

Compact sets are generally defined on *metric spaces*. What is this?

## Definition 1.1: Metric Function

Let  $X$  be a nonempty set. A function  $d : X \times X \rightarrow \mathbb{R}$  is called a **metric function** on  $X$  if

- $d(x, y) > 0$  for  $\forall x, y \in X$  with  $x \neq y$
- $d(x, y) = 0 \Leftrightarrow x = y$
- $d(x, y) = d(y, x)$  for  $\forall x, y \in X$
- $d(x, y) \leq d(x, z) + d(z, y)$  for  $\forall x, y, z \in X$ .

## Definition 1.2: Metric Space

A nonempty set  $X$  with a metric function  $d : X \times X \rightarrow \mathbb{R}$  is called a metric space.

**Example 1**

$\mathbb{R}$  is a metric space with a metric function  $d(x, y) = |x - y|$ . We claim that the metric  $d(x, y) = |x - y|$  satisfies the four axioms of a metric function.

- If  $x, y \in \mathbb{R}$  with  $x \neq y$ , then  $|x - y| > 0$ .
- $|x - x| = 0$ , and if  $|x - y| = 0$  then  $x = y$
- For all  $x, y \in \mathbb{R}$ ,  $|x - y| = |y - x|$ .
- For all  $x, y, z \in \mathbb{R}$ ,  $|x - y| \leq |x - z| + |z - y|$  (triangle inequality).

Therefore,  $\mathbb{R}$  with  $d(x, y) = |x - y|$  is a metric space.

**Example 2**

Let  $X$  be the set of all bounded real-valued functions on  $A (\neq \emptyset)$ . For  $f, g \in X$ , we define  $d(f, g) = \sup \{|f(x) - g(x)| : x \in A\}$ . Since

- $0 \leq |f(x) - g(x)| \leq |f(x)| + |g(x)| \leq 2M$  for all  $x \in A$
- $d(f, g) = 0 \Leftrightarrow f = g$  since  $|f(x) - g(x)| \leq d(f, g)$  for  $\forall x \in A$
- $d(f, g) = d(g, f)$
- $\sup\{|f(x) - g(x)| : x \in A\} = \sup\{|f(x) - h(x) + h(x) - g(x)| : x \in A\} \leq \sup\{|f(x) - h(x)| : x \in A\} + \sup\{|h(x) - g(x)| : x \in A\} = d(f, h) + d(h, g),$

$d$  is a metric on  $X$ .

With this definition, we can generalize what we have done on  $\mathbb{R}$  to a metric space.

**Definition 1.3: Neighborhood**

Let  $(X, d)$  be a metric space. For  $\epsilon > 0$  and  $p \in X$ , the set

$$N_\epsilon(p) = \{x \in X \mid d(p, x) < \epsilon\}$$

is called an  $\epsilon$ -**neighborhood** of  $p$ .

## 2

## Open and Closed Sets

From now on, denote  $X$  as a metric space. The proofs for results are omitted if they have the exact same proof with the case  $X = \mathbb{R}$ .

**Definition 2.1: Interior Point**

Let  $E \subset X$  be a set. A point  $p$  is an **interior point** of  $E$  if for some  $\epsilon > 0$ , the  $\epsilon$ -neighborhood of  $p$  is entirely in  $E$ . That is,  $N_\epsilon(p) \subset E$ .

**Definition 2.2: Isolated Point**

Let  $E \subset X$  be a set. A point  $p$  is an **isolated point** of  $E$  if for some  $\epsilon > 0$ ,  $p$  is the only point of  $E$  in the  $\epsilon$ -neighborhood of  $p$ . That is,  $N_\epsilon(p) \cap E = \{p\}$ .

**Definition 2.3: Boundary Point**

Let  $E \subset X$  be a set. A point  $p$  is a **boundary point** of  $E$  if any neighborhood of  $c$  contains points from both  $E$  and  $E^c$ . That is,  $N_\epsilon(p) \cap E \neq \emptyset$  and  $N_\epsilon(p) \cap E^c \neq \emptyset$ .

**Definition 2.4: Limit Point**

Let  $E \subset X$  be a set. A point  $p$  is a **limit point** of  $E$  if any  $\epsilon$ -neighborhood of  $p$  contains a point of  $E$  other than  $p$ . That is,  $N_\epsilon(p) \cap (E \setminus \{p\}) \neq \emptyset$ .

**Proposition.**

$p$  is a limit point of  $E$  if and only if there is a sequence of elements in  $E \setminus \{p\}$  converging to  $p$ .

*Proof.* ( $\Leftarrow$ ) Suppose the sequence  $\{x_n\}$  of elements of  $E \setminus \{p\}$  converges to  $p$ . Then  $\forall \epsilon > 0$ ,  $\exists N$  such that  $\forall n \geq N$ ,  $d(x_n, p) < \epsilon$ . Thus  $p$  is a limit point of  $E$ .

( $\Rightarrow$ ) If  $p$  is a limit point of  $E$ , choose  $x_n \in E \setminus \{p\}$  with  $d(x_n, p) < 1/n$ , then  $x_n \rightarrow p$ . ■

**Definition 2.5: Open and Closed Sets**

A set  $S \subset \mathbb{R}$  is **open** if every point of  $S$  is an interior point. A set  $S \subset \mathbb{R}$  is **closed** if it contains all of its limit points.

Note that the definitions coincide to the definitions discussed in class if we set  $X = \mathbb{R}$ .

**Theorem 2.1**

- A complement of an open set is closed.
- A complement of a closed set is open.

**Theorem 2.2**

Let  $(X, d)$  be a metric space.

1. If  $\{O_\alpha\}_{\alpha \in A}$  is a collection of open sets of  $X$ , then  $\bigcup_{\alpha \in A} O_\alpha$  is open. That is, an arbitrary union of open sets is open.
2. If  $\{O_1, \dots, O_n\}$  is a finite collection of open sets of  $X$ , then  $\bigcap_{j=1}^n O_j$  is open. That is, a finite intersection of open sets is open.

*Proof.* (1) We may assume that  $\bigcup_{\alpha \in A} O_\alpha \neq \emptyset$ . Let  $p \in \bigcup_{\alpha \in A} O_\alpha$ , then  $p \in O_\alpha$  for some  $\alpha \in A$ . Since  $O_\alpha$  is open,  $\exists \epsilon > 0$  such that  $N_\epsilon(p) \subseteq O_\alpha \subseteq \bigcup_{\alpha \in A} O_\alpha$ . Thus,  $p$  is an interior point of  $\bigcup_{\alpha \in A} O_\alpha$ .

(2) Assume  $\bigcap_{j=1}^n O_j \neq \emptyset$ . Let  $p \in \bigcap_{j=1}^n O_j$ . Then  $p \in O_j$  for  $\forall j = 1, 2, \dots, n$ . Since each  $O_j$  is open,  $\exists \epsilon_j > 0$  such that  $N_{\epsilon_j}(p) \subseteq O_j$ . Now, let  $\epsilon = \min\{\epsilon_1, \epsilon_2, \dots, \epsilon_n\} > 0$ , then  $N_\epsilon(p) \subseteq N_{\epsilon_j}(p) \subseteq O_j$  for all  $j$ . Therefore,  $N_\epsilon(p) \subseteq \bigcap_{j=1}^n O_j$ , and  $p$  is an interior point. ■

**Corollary**

Let  $(X, d)$  be a metric space.

1. If  $\{O_1, \dots, O_n\}$  is a finite collection of closed sets of  $X$ , then  $\bigcup_{j=1}^n O_j$  is closed. That is, a finite union of closed sets is closed.
2. If  $\{O_\alpha\}_{\alpha \in A}$  is a collection of closed sets of  $X$ , then  $\bigcap_{\alpha \in A} O_\alpha$  is closed. That is, an arbitrary intersection of open sets is open.

## 3

## Compact Sets

Now, we state the general definition of compact sets.

**Definition 3.1: Open Cover**

Let  $X$  be a metric space, and  $E \subset X$ . A collection  $\{O_\alpha\}_{\alpha \in A}$  of open subsets of  $X$  is an **open cover** of  $E$  if

$$E \subseteq \bigcup_{\alpha \in A} O_\alpha.$$

**Definition 3.2: Compact Set**

Let  $X$  be a metric space. A set  $K \subseteq X$  is **compact** if every open cover of  $K$  has a finite subcover of  $K$ .

That is, if  $\{O_\alpha\}$  is an open cover of  $K$ ,  $K$  is compact if  $\exists \alpha_1, \dots, \alpha_n \in A$  such that

$$K \subseteq \bigcup_{j=1}^n O_{\alpha_j}.$$

**Example 3**

Every finite set is compact.

**Example 4**

$I = (0, 1)$  is not compact. Consider  $O_n = (0, \frac{n}{n+1})$  for  $n \in \mathbb{N}$ . Then,  $\{O_n\}_{n \in \mathbb{N}}$  is an open cover of  $I$ . Indeed, if  $x \in I$ , then  $\exists n_0 \in \mathbb{N}$  such that  $\frac{1}{n_0+1} < 1 - x$  by the Archedian property. Thus,

$$x \in O_{n_0} \subseteq \bigcup_{n=1}^{\infty} O_n.$$

But, no finite subcover can cover  $I$ . Assume to the contrary that a finite subcover  $\{O_{n_1}, O_{n_2}, \dots, O_{n_k}\}$  covers  $I$ . Let  $N = \max\{n_1, \dots, n_k\}$ , then we have

$$(0, 1) \subseteq \bigcup_{j=1}^k O_{n_j} = \left(0, \frac{N}{N+1}\right),$$

which gives a contradiction.

## 4

## Properties of Compact Sets

**Theorem 4.1: Heine-Borel**

Every closed and bounded interval  $[a, b]$  is compact.

*Proof.* Let  $\mathcal{U} = \{U_\alpha\}_{\alpha \in A}$  be an open cover of  $[a, b]$ . Define

$$E = \{r \in [a, b], [a, r] \text{ is covered by a finite subcover of } \mathcal{U}\}.$$

Clearly,  $E$  is nonempty and bounded. Thus  $\exists \gamma = \sup E$  in  $\mathbb{R}$  by the least upper bound property.

**Claim.**  $\gamma = b$ .

Suppose that  $\gamma < b$ . We will find a contradiction by constructing  $s \in E$  such that  $\gamma < s$ . Since  $\gamma \in U_\alpha$  for some open set  $U_\alpha \in \mathcal{U}$ ,  $\exists \epsilon > 0$  such that  $N_\epsilon(\gamma) = (\gamma - \epsilon, \gamma + \epsilon) \subseteq U_\alpha$ . Since  $\gamma - \epsilon$  is not an upper bound of  $E$ ,  $\exists t \in E$  such that  $\gamma - \epsilon < t < \gamma$ . Thus,  $[0, t]$  is covered by finitely many sets

$$U_{\alpha_1}, U_{\alpha_2}, \dots, U_{\alpha_n}.$$

Now, choose any  $s \in (\gamma, \gamma + \epsilon)$  such that  $s < b$ . Then,

$$[a, s] \subseteq \left( \bigcup_{j=1}^n U_{\alpha_j} \right) \cup U_\alpha,$$

i.e.  $s \in E$ . Also since  $\gamma \in E$ , so this completes the proof. ■

Note that if  $X = \mathbb{R}$ , closed and bounded is equivalent to compact (so a compact set is also closed and bounded). In a general metric space, every closed and bounded set is compact, but not every compact set is closed and bounded.

Then, is this definition equivalent to the sequential definition for  $X = \mathbb{R}$ ? Yes!

**Theorem 4.2**

Let  $K \subset \mathbb{R}$ . Then  $K$  is compact if and only if every sequence in  $K$  has a subsequence that converges to a point in  $K$ .

*Proof.* ( $\Rightarrow$ ) Let  $\{p_n\}$  be a sequence in  $K$ , and let  $E = \{p_n \mid n = 1, 2, \dots\}$ . If  $E$  is finite, then there exists a point  $p \in E$  and a sequence  $\{n_k\}$  with  $n_1 < n_2 < \dots$  such that

$$p_{n_1} = p_{n_2} = \dots = p.$$

The sequence  $\{p_{n_k}\}$  obviously converges to  $p \in K$ .

If  $E$  is infinite, then  $E$  has a limit point  $p \in K$ . Choose  $n_1$  such that  $d(p, p_{n_1}) < 1$ . Having chosen  $n_1, \dots, n_{k-1}$ , choose an integer  $n_k > n_{k-1}$  so that

$$d(p, p_{n_k}) < \frac{1}{k}.$$

Such an integer  $n_k$  exists since every neighborhood of  $p$  contains infinitely many points of  $E$ . The sequence  $\{p_{n_k}\}$  is a subsequence of  $\{p_n\}$  converging to  $p \in K$ .

( $\Leftarrow$ ): Let  $p$  be a limit point of  $K$ . Then there exists a sequence of distinct points in  $K$  converging to  $p$ . Since each of its subsequence converge to  $p$ , hence  $p \in K$ , so  $K$  is closed.

Assume  $K$  is not bounded. For each  $k \in \mathbb{N}$ , choose a point  $p_k \in K$  such that  $d(p_k, p_0) \geq k$  for some fixed  $p_0 \in X$ . Then the sequence  $\{p_k\}$  satisfies  $d(p_k, p_0) \rightarrow \infty$ , so it cannot have any convergent subsequence (since convergent sequences are bounded), a contradiction.

Since  $K$  is closed and bounded, it is compact. ■

#### Theorem 4.3

Let  $X$  be a metric space. If  $K \subseteq X$  is compact, then

1.  $K$  is closed
2. If  $F \subseteq K$  and  $F$  is closed, then  $F$  is compact.

*Proof.* (1) It is enough to show that  $K^\complement$  is open. Let  $p \in K^\complement$ . For each  $q \in K$ , Let  $\epsilon_q = d(p, q)/2$ . Then,  $N_{\epsilon_q}(p) \cap N_{\epsilon_q}(q) = \emptyset$ . Since  $\{N_{\epsilon_q}(q)\}_{q \in K}$  is an open cover of  $K$ , there exists  $q_1, q_2, \dots, q_n$  such that

$$K \subseteq \bigcup_{j=1}^n N_{\epsilon_{q_j}}(q_j).$$

Let  $\epsilon = \min\{\epsilon_1, \dots, \epsilon_n\}$ . Then,  $N_\epsilon(p)$  does not intersect with  $N_{\epsilon_{q_j}}(q_j)$  for all  $j = 1, \dots, n$ . Thus,  $N_\epsilon(p) \subseteq K^\complement$ , which proves that  $K$  is closed.

(2) Let  $\{O_\alpha\}_{\alpha \in A}$  be an open cover of  $F$ . Then,

$$\{O_\alpha\}_{\alpha \in A} \cup F^\complement$$

is an open cover of  $K$ . Since  $K$  is compact,  $\exists$  a finite subcollection of  $\{O_\alpha\}_{\alpha \in A} \cup F^\complement$  containing  $K$ , which also contains  $F$ . ■

#### Corollary

If  $F$  is closed and  $K$  is compact, then  $F \cap K$  is compact.

The previous theorem gives a simple proof for generalized Cantor's intersection property.

**Corollary : Cantor's Intersection Property**

If  $K_1 \supset K_2 \supset \dots$  is a nested family of nonempty compact sets, then their intersection  $K = \bigcap_{n=1}^{\infty} K_n$  is nonempty and compact.

*Proof.* Since  $K$  is a closed subset of a compact set  $K_n$ , it is compact. ■

**Theorem 4.4**

The continuous image of a compact set is compact.

*Proof.* Let  $f : A \rightarrow f(A)$  be a continuous function on a compact set  $A$ . For any sequence  $\{y_n\}$ , we can find a corresponding sequence  $\{x_n\}$  such that  $f(x_n) = y_n$  for all  $n \in \mathbb{N}$ . Since there exist a convergent subsequence  $\{x_{n_k}\}$  converging to  $x \in A$ , the subsequence  $\{y_{n_k}\}$  converges to  $y = f(x) \in f(A)$ . Therefore, every sequence in  $f(A)$  has a subsequence converging to a point in  $f(A)$ , and  $f(A)$  is compact. ■

**Remark.**

If  $X = \mathbb{R}$ , then the continuous image of a closed set need not be closed.

**Remark.**

If  $X = \mathbb{R}$ , then the continuous image of a bounded set need not be bounded.

This gives the Extreme value theorem.

**Corollary : Extreme Value Theorem**

Let  $K \in \mathbb{R}$  be a compact set. If  $f : K \rightarrow \mathbb{R}$ , then  $f(x)$  attains its minimum and maximum value on  $K$ .